ILLINOIS BONE & JOINT INSTITUTE®

MEDICATION INSTRUCTIONS

Jeffrey M. Goldstein, MD

9000 Waukegan Road, Suite 200 ~ Morton Grove, IL 60053 (847) 375-3000

Patient:	MR#: .30*	#15

Because of the way certain medications work within your body and interact with each other, it may be important to stop taking them prior to your joint replacement surgery.

Please check all the medications you are currently taking and identify them on the list I have compiled. This list will assist you in determining if and when you need to stop your medications. If your medication is not listed or you have been specifically instructed otherwise, you may continue to take your medication, as usual, until the day before surgery.

Please notify me if you are taking narcotic pain medication on a regular basis. Weaning off or stopping the medications listed below is highly recommended prior to surgery:

Hydrocodone

Norco

Oxycodone

Oxycontin

Percocet

Vicodin

One (1) week or 7 days prior to surgery, please stop taking the following medications:

All birth control pills & any male or female hormone (including creams and patches) such as:

Emcyt, Estraderm, Estratest, Estrace, Estradiol, Estrogens, Ogen, Premarin, Prempro,
Testosterone.

All vitamins such as: A, C, E, K, Multivitamins, Fish Oil, Omega (3,6,9), Juice Plus, CoQ10

All herbal supplements such as: Echinacea, Ephedra, Garlic, Ginkgo, Ginseng, Kava, St. John's Wort, Valerian, Saw Palmetto

All joint supplements such as: Glucosamine, Chondroitin, MSM

All Anti-Coagulants such as: Coumadin, Warfarin, Xarelto, Pradaxa

All Anti-Platelets such as: Plavix, Clopidogrel, Aggrenox, Dipyridamole, Pletal, Cilostazol, Trental, Pentoxil

NOTE: If you have heart stents, do not stop Plavix until seen by a cardiologist prior to surgery.

All Aspirin containing products such as: Anacin, Ascriptin, Bayer, Bufferin, Ecotrin, Excedrin, Fasprin, Norgesic, Percodan

NOTE: If you have heart stents and take Plavix and Aspirin, do not stop Aspirin

All Anti-Thrombotics such as: Ticlid, Ticlodipine

MEDICATION INSTRUCTIONS - Continued

Choline and Magnesium Salicylates (CMT, Tricosal, Trilisate)

Choline Salicylate (Arthropan)

Diclofenac (Cataflam, Voltaren, Arthrotec)

Diflunisal (Dolobid)

Etodolac (Lodine)

Fenoprofen Calcium (Nalfon)

Flurbiprofen (Ansaid)

Ibuprofen (Advil, Motrin, Nuprin)

Indomethacin (Indocin)

Ketoprofen (Actron, Orudis, Oruvail)

Magnesium Salicylate (Arthritab, Bayer Select, Doan's Pills, Magan, Mobidin, Mobogesic)

Meloxicam (Mobic)

Nabumetone (Relafen)

Naproxen (Aleve, Naprosyn, Naprelan, Anaprox)

Oxaprozin (Daypro)

Piroxicam (Feldene)

Salsalate (Amigesic, Anaflex 750, Disalcid, Marthritic, Mono-Gesic, Salfex, Salsitab)

Sodium Salicylate (various generics)

Sulindac (Clinoril)

Two (2) days prior to surgery, please stop taking the following medications:

Benazepril (Lotensin)

Candesartan (Atacand)

Enalapril (Vasotec, Renitec)

Eprosartan (Teveten)

Fosinopril (Monopril)

Irbesartan (Avapro)

Lisinopril (Lisodur, Lopril, Novatec, Prinival, Zestril)

Losartan (Cozaar)

Olmesartan (Benicar)

Perindopril (Coversy, Aceon)

Quinapril (Accupril)

Ramipril (Altrace, Tritace, Ramace, Ramiwin)

Telmisartan (Micardis)

Valsartan (Diovan)

Zofenopril

You may continue to take the following medications until the time of surgery:

Acetaminophen

Celebrex

Hydrochlorothiazide

Tramadol

Tylenol

MEDICATION INSTRUCTIONS – Continued

You may continue to take the following medications until the time of surgery:

Beta Blockers:

Acebutolol

Atenolol (Tenormin)

Bisoprolol (Zabeta, Ziac)

Coreg (Carvedilol)

Corgard (Nadolol)

Inderal (Propranolol)

Labetolol

Metoprolol (Toprol)

Pindolol

Calcium Channel Blockers:

Amlodipine (Norvasc)

Cardizem (Diltiazem)

Nifedipine (Procardia)

Verapamil

Contact your medical doctor for instructions if you take any of the following medications:

Diabetic Medications Containing Metformin:

ACTOplus met

Avandamet

Glucovance

Janumet

Metaglip

PrandiMet

Adderall

Cytoxan

Enbril

Imuran

Librax

Librium

Methotrexate

Remicade

I encourage you to contact me as soon as possible if you have questions related to your medications.